

Name of research institute or organization:

**Institut d'Astrophysique et de Géophysique, Université de Liège**

Title of project:

High resolution, solar infrared Fourier Transform Spectrometry. Application to the study of the Earth atmosphere

Project leader and team:

Luc Delbouille (em.), Philippe Demoulin, Pierre Duchatelet, Emmanuel Mahieu, Ginette Roland (em.), Christian Servais (project leader), Rodolphe Zander (em.)  
Jacqueline Bosselois, Guy Buntinx, Olivier Flock, Vincent Van De Weerd, Diane Zander

Project description:

The main activity of the Liège group at the Jungfraujoch was the continuation of the long-term monitoring of the Earth atmosphere. The observations achieved by the two high-performance infrared spectrometers allow to routinely derive abundances of more than 20 constituents related to the erosion of the ozone layer in the stratosphere (HCl, ClONO<sub>2</sub>, HNO<sub>3</sub>, NO, NO<sub>2</sub>, HF, COF<sub>2</sub>, O<sub>3</sub>...), monitored in the frame of the Kyoto protocol (N<sub>2</sub>O, CH<sub>4</sub>, CO<sub>2</sub>, SF<sub>6</sub>, CCl<sub>2</sub>F<sub>2</sub>, CHClF<sub>2</sub>, CCl<sub>3</sub>F...) or affecting the oxidization processes in the troposphere (CO, C<sub>2</sub>H<sub>2</sub>, C<sub>2</sub>H<sub>6</sub>, OCS, HCN, H<sub>2</sub>CO...). The resulting databases allow the determination of the short-term variability, seasonal modulations, as well as long-term changes affecting most of these species.


During 2005, Liège observers spent 247 days at the Jungfraujoch. Good weather conditions enabled observations on 124 days.

For a number of the species listed above, a complete re-analysis of the archived spectra is currently under way with SFIT-2, a recent retrieval algorithm that provides in most cases information on the distribution of the molecules versus altitude. This algorithm allows determining partial columns (e.g. to distinguish between tropospheric and stratospheric contents) as well as more accurate total columns.

In the frame of the EC project UFTIR (<http://www.nilu.no/uftir>), a homogenised optimal retrieval strategy has been developed for the inversion of O<sub>3</sub>, N<sub>2</sub>O, C<sub>2</sub>H<sub>6</sub>, HCFC-22, CO and CH<sub>4</sub>. The corresponding Jungfraujoch spectra between 1995 and 2004 have been reprocessed, after implementation of this new strategy, and the resulting time series are being archived on the UFTIR database at NILU. The UFTIR outcomes are shared with the NDSC infrared community.

As an example, Figure 1 shows the results of the FTIR retrievals for carbon monoxide CO, compared to the Jungfraujoch *in situ* CO measurements by EMPA.


The agreement between both datasets is rather good, showing similar seasonal and short term variations. CO anomalies observed in the northern hemisphere in 1998-1999, 2002 and 2003 are clearly visible. These anomalies have recently been investigated [Yurganov *et al.*, 2005] and the corresponding extra CO emissions have been evaluated to 95 and 130 Tg, respectively in 2002 and 2003. Strong boreal fires that occurred in Russia during these two years are the most likely causes for the observed CO burden increases.



**Figure 1:** Daily mean carbon monoxide volume mixing ratio (VMR) at the Jungfraujoch, derived from FTIR spectra (circles) and from EMPA *in situ* measurements (triangles). FTIR data correspond to the mean VMR retrieved in the lowest layers (i.e. between 3.58 and 5.5 km, corresponding to one independent piece of information). EMPA data have been obtained from the WDCGG (World Data Centre for Greenhouse Gases, <http://gaw.kishou.go.jp/wdcgg.html>).

Within the context of the Montreal Protocol, monitoring of chlorinated source gases has been part of our activities. Time series of CFC-12 ( $\text{CCl}_2\text{F}_2$ ) and HCFC-22 ( $\text{CHClF}_2$ ) have been updated while modifications implemented to the SFIT-2 retrieval algorithm have allowed to perform retrievals of two additional species, i.e. CFC-11 ( $\text{CCl}_3\text{F}$ ) and  $\text{CCl}_4$ . Current monthly mean time series of three of these source gases are shown in Figure 2. Although year round data points are reproduced here, only averaged total columns of the quietest June to November months (i.e. exhibiting less variability, see filled symbols) have been used to characterise the long-term evolutions of these compounds. Corresponding trends and annual column changes are available in Zander *et al.* [2005]. It is interesting to point out here the contrasted evolutions of these source gases: (i) the Montreal-controlled CFC-11 and CFC-12 are decreasing/stabilising as expected from their respective lifetimes (45 and 100 years); (ii) progressive phase out of HCFC-22 (an important CFC substitute) has begun in 2004 and its production is supposed to reach zero in developed countries in 2030; its concentration is expected to continue rising until about 2010 before stabilisation and rapid decrease thereafter. Our measurements indicate a steady increase of the HCFC-22 total columns with recent rate of change on the order of 3 %/year.

Comparisons of the above results with findings deduced from *in situ* measurements performed by the AGAGE and NOAA/CMDL networks indicate very good agreement in terms of trends and atmospheric concentrations for these various species.



**Figure 2:** Contrasted evolutions of the monthly mean total vertical column abundances of CFC-12, CFC-11 and HCFC-22 above the Jungfrauoch.

In addition to the important greenhouse gases regulated by the Montreal Protocol, the 1997 "Kyoto Protocol on climate change" specifically targets  $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{N}_2\text{O}$  and  $\text{SF}_6$  which present characteristic infrared absorption features allowing to quantify their atmospheric abundances. Regular analyses of all Jungfrauoch observations have resulted in updates of their temporal evolutions [Zander *et al.*, 2005]. Related time series, which cover now two decades, are reproduced in Figure 3.

The first obvious feature is the regular growth for three of these gases over the 1985-2004 time period, only methane exhibiting a stabilisation over recent years.


Comparisons with abundances derived from pioneering observations performed in 1950-1951 at the same site by M. Migeotte indicate that the total columns of  $\text{CO}_2$ ,  $\text{CH}_4$  and  $\text{N}_2\text{O}$  have been respectively multiplied by 1.25, 1.35 and 1.17. More specifically, trend determinations have indicated a yearly increase of 0.42 % for  $\text{CO}_2$ , in excellent agreement with *in situ* surface measurements (e.g. [www.cmdl.noaa.gov](http://www.cmdl.noaa.gov)). The very long-lived nitrous oxide (120 years) shows a similar behaviour, with an annual linear build up of  $1.06 \times 10^{16}$  molec./ $\text{cm}^2$ ; this corresponds to an increase of 0.26 %/year, commensurate with *in situ* trend data.

Rapid increase of the total column abundance of  $\text{SF}_6$  has been confirmed over recent years, with an annual load increase still exceeding 4 %/year in 2004. It is important to limit emissions of this compound to the atmosphere because it combines a very strong

absorption of infrared radiation on a per-molecule basis with a very long lifetime of several thousand years.

Extrapolation of the Jungfraujoch data predicts tropospheric SF<sub>6</sub> concentrations of about 15 pptv in 2050 and about 25 pptv in 2100 (compared this to the 2.0 pptv concentration measured in 1988) [Krieg *et al.*, 2005]. These values are significantly lower than those reported in recent scenarios [WMO 2003], justifying future monitoring of this species to determine its effective future evolution and related climatic impact.

A striking feature of this figure is the stabilisation of the CH<sub>4</sub> loading during recent years, that will deserve future comparisons with models, to identify the relative contributions of changes in sources and sinks leading to this stabilisation.



**Figure 3:** Long-term evolution of four species targeted by the Kyoto Protocol as derived from ground-based remote observations conducted at the Jungfraujoch station. Notice the different vertical axis units for each frame.

During 2005, we provided additional data for the calibration/validation of 3 instruments (MIPAS, SCIAMACHY and GOMOS) aboard the European satellite Envisat. On the whole, we supplied to the calibration team 12784 total column abundances of O<sub>3</sub>, N<sub>2</sub>O, CO, CH<sub>4</sub>, NO, NO<sub>2</sub>, HNO<sub>3</sub> and CO<sub>2</sub>, deduced from Jungfraujoch observations between July 2002 and December 2004. More elaborated products consisting in 4329 vertical distributions and related partial columns of HNO<sub>3</sub>, CH<sub>4</sub> and N<sub>2</sub>O for 201 days between July 2002 and March 2004 were also produced for specific validation of MIPAS profiles and SCIAMACHY columns.

The Canadian ACE-FTS instrument was launched in August 2003 and has been in "post commissioning" operation since February 2004 [Bernath *et al.*, 2005]. Specific observational campaigns were organized at the Jungfraujoch in support to the

validation of the ACE-FTS spectrometer to record as many coincident measurements as possible.

Scientific and validation papers using the first "Version 1" of level 2 data have been published recently in a GRL special issue. In particular, a study dealing with comparisons between stratospheric columns of HCl and ClONO<sub>2</sub> measured up to October 2004 by ACE and by ground-based FTIR instruments operated at five northern latitude NDSC sites (including the Jungfraujoch) has been led by ULg [Mahieu et al., 2005]. Main conclusions were that: (i) ACE is able to identify for both targeted gases distribution features characteristic of geographical, dynamical, seasonal and chemical changes occurring in the atmosphere; (ii) excellent agreement was found when considering the only sets of coincident measurements obtained around the Thule site (76.5°N), with mean partial column ratio (ACE/Thule) equal to 1.04 and 0.99 respectively for HCl and ClONO<sub>2</sub>; (iii) good agreement is generally found for other sites (in particular for the Jungfraujoch), even if systematic differences between some data sets deserve further investigations based on additional coincident ACE "Version 2.2" and ground-level remote FTIR measurements.

On the hardware side, the opening of the heliostat has been electrified, a first step towards its complete automation. The previously tedious manual opening of the heliostat is now performed with a simple remote control.

Key words:

---

Earth atmosphere, ozone layer, greenhouse gases, long-term monitoring, infrared spectroscopy

Internet data bases:

---

<http://www.nilu.no/nadir/>, <ftp://ndsc.ncep.noaa.gov/pub/ndsc/jungfrau/ftir/>

Collaborating partners/networks:

---

Main collaborations: IASB (Institut d'Aéronomie Spatiale de Belgique) / NDSC (Network for the Detection of Stratospheric Change) / SOGE partners (e.g. EMPA) [<http://www.nilu.no/soge/>] / NASA Langley Research Center / NASA JPL / University of Oslo / University of Leeds / IMK (Forschungszentrum Karlsruhe) / satellite experiments: MOPPIT, ENVISAT and ACE validation / ...

Scientific publications and public outreach 2005:

---

### Refereed journal articles

Barret, B., D. Hurtmans, M.R. Carleer, M. De Mazière, E. Mahieu, and P.-F. Coheur, Line narrowing effect on the retrieval of HF and HCl vertical profiles from ground-based FTIR measurements, *J. Quant. Spectrosc. Radiat. Transfer*, **95**, 499-519, 2005.

Bernath, P.F., C.T. McElroy, M.C. Abrams, C.D. Boone, M. Buttler, C. Camy-Peyret, M. Carleer, C. Clerbaux, P.-F. Coheur, R. Colin, P. DeCola, M. De Mazière, J.R. Drummond, D. Dufour, W.F.J. Evans, H. Fast, D. Fussen, K. Gilbert, D.E. Jennings, E.J. Llewellyn, R.P. Lowe, E. Mahieu, J.C. McConnell, M. McHugh, S.D. McLeod, R. Michaud, C. Midwinter, R. Nassar, F. Nichitiu, C. Nowlan, C.P. Rinsland, Y.J. Rochon, N. Rowlands, K. Semeniuk, P. Simon, R. Skelton, J.J. Sloan, M.-A. Soucy, K. Strong, P. Tremblay, D. Turnbull, K.A. Walker, I. Walkty, D.A. Wardle, V. Wehrle, R. Zander, and J. Zou, Atmospheric Chemistry Experiment (ACE): mission overview, *Geophys. Res. Lett.*, **32**, L15S01, doi:10.1029/2005GL022386, 2005.

De Mazière, M., C. Vigouroux, T. Gardiner, M. Coleman, P. Woods, K. Ellingsen, M. Gauss, I. Isaksen, T. Blumenstock, F. Hase, I. Kramer, C. Camy-Peyret, P. Chelin, E. Mahieu, P. Demoulin, P. Duchatelet, J. Mellqvist, A. Strandberg, V. Velazco, J. Notholt, R. Sussmann, W. Stremme, and A. Rockmann, The exploitation of ground-based Fourier transform infrared observations for the evaluation of tropospheric trends of greenhouse gases over Europe, *Environmental Sciences*, **2** (2-3), 283-293, June-September 2005.

Krieg, J., J. Notholt, E. Mahieu, C.P. Rinsland, and R. Zander, Sulphur hexafluoride (SF<sub>6</sub>): comparison of FTIR-measurements at three sites and determination of its trend in the northern hemisphere, *J. Quant. Spectrosc. Radiat. Transfer*, **92**, 383-392, 2005.

Mahieu, E., R. Zander, P. Duchatelet, J.W. Hannigan, M.T. Coffey, S. Mikuteit, F. Hase, T. Blumenstock, A. Wiacek, K. Strong, J.R. Taylor, R. Mittermeier, H. Fast, C.D. Boone, S.D. McLeod, K.A. Walker, P.F. Bernath, and C.P. Rinsland, Comparisons between ACE-FTS and ground-based measurements of stratospheric HCl and ClONO<sub>2</sub> loadings at northern latitudes, *Geophys. Res. Lett.*, **32**, L15S08, doi:10.1029/2005GL022396, 2005.

Rinsland, C.P., C. Boone, R. Nassar, K. Walker, P. Bernath, E. Mahieu, R. Zander, J.C. McConnell, and L. Chiou, Trends of HF, HCl, CCl<sub>2</sub>F<sub>2</sub>, CCl<sub>3</sub>F, CHClF<sub>2</sub> (HCFC-22), and SF<sub>6</sub> in the lower stratosphere from Atmospheric Chemistry Experiment (ACE) and Atmospheric Trace MOlecule Spectroscopy (ATMOS) measurements near 30°N latitude, *Geophys. Res. Lett.*, **32**, L16S03, doi:10.1029/2005GL022415, 2005.

Rinsland, C.P., A. Goldman, E. Mahieu, R. Zander, L.S. Chiou, J.W. Hannigan, S.W. Wood, and J.W. Elkins, Long-term evolution in the tropospheric concentration of chlorofluorocarbon 12 (CCl<sub>2</sub>F<sub>2</sub>) derived from high-spectral resolution infrared solar absorption spectra: retrieval and comparison with *in situ* surface measurements, *J. Quant. Spectrosc. Radiat. Transfer*, **92**, 201-209, 2005.

Yurganov, L.N., P. Duchatelet, A.V. Dzhola, D.P. Edwards, F. Hase, I. Kramer, E. Mahieu, J. Mellqvist, J. Notholt, P.C. Novelli, A. Rockmann, H.E. Scheel, M. Schneider, A. Schulz, A. Strandberg, R. Sussmann, H. Tanimoto, V. Velazco, J.R. Drummond, and J.C. Gille, Increased Northern Hemispheric carbon monoxide burden in the troposphere in 2002 and 2003 detected from the ground and from space, *Atmos. Chem. Phys.*, **5**, 563-573, 2005.

Zander, R., E. Mahieu, P. Demoulin, P. Duchatelet, C. Servais, G. Roland, L. Delbouille, M. De Mazière and C.P. Rinsland, Evolution of a dozen non-CO<sub>2</sub> greenhouse gases above Central Europe since the mid-1980s, *Environmental Sciences*, **2** (2-3), 295-303, June-September 2005.

### Conference papers

Mahieu, E., R. Zander, P. Demoulin, P. Duchatelet, C. Servais, C.P. Rinsland, and M. De Mazière, Recent evolution of atmospheric OCS above the Jungfraujoch station: Implications for the stratospheric aerosol layer, in *Proceedings of "Atmospheric Spectroscopy Applications, ASA Reims 2005"*, Reims, September 6-8, 2005, pp.235-238, 2005.

### Magazine and Newspapers articles

"La couche d'ozone se reconstruit. Des chercheurs de l'ULg étudient la composition chimique de l'atmosphère depuis un sommet suisse", with Pierre Duchatelet, Groupe Sud Presse, 17 March 2005.

### Radio and television

"La sentinelle du soleil", interview of Pierre Duchatelet, Belgian Local Television No Télé (Hainaut), 19 October 2004.

Address:

---

Institut d'Astrophysique et de Géophysique - Université de Liège  
allée du VI août, 17 - Bâtiment B5a  
B-4000 Sart Tilman (Liège, Belgique)

Contacts:

---

|                   |                     |                                    |
|-------------------|---------------------|------------------------------------|
| Luc Delbouille    | Tel. +32 4 342 2594 | e-mail: delbouille@astro.ulg.ac.be |
| Philippe Demoulin | Tel. +32 4 366 9785 | e-mail: demoulin@astro.ulg.ac.be   |
| Pierre Duchatelet | Tel. +32 4 366 9786 | e-mail: duchatelet@astro.ulg.ac.be |
| Emmanuel Mahieu   | Tel. +32 4 366 9786 | e-mail: mahieu@astro.ulg.ac.be     |
| Ginette Roland    | Tel. +32 4 342 2594 | e-mail: roland@astro.ulg.ac.be     |
| Christian Servais | Tel. +32 4 366 9784 | e-mail: servais@astro.ulg.ac.be    |
| Rodolphe Zander   | Tel. +32 4 366 9756 | e-mail: zander@astro.ulg.ac.be     |

URL: <http://girpas.astro.ulg.ac.be/>

