

Name of research institute or organization:

**Bundesamt für Strahlenschutz, Freiburg i.Br.
Climate and Environmental Physics, University of Bern**

Title of project:

⁸⁵Kr Activity Determination in Tropospheric Air

Project leader and team

Hartmut Sartorius, Clemens Schlosser and Sabine Schmid, Bundesamt für Strahlenschutz, D-79098 Freiburg

Roland Purtschert, Heinz Hugo Loosli, Physikalisches Institut, Universität Bern, CH-3012 Bern

Project description:

The collection of air samples for ⁸⁵Kr activity measurements has been continued at Jungfrauoch in 2005. A few cc of Krypton are collected in weekly samples from about 10 m³ of air. These samples are sent to Freiburg i.Br. for Krypton separation, purification and for activity measurement.

This isotope is unique because it contributes the major part to the present-day artificial activity in air. The radiation dose however is negligible compared to the dose components from internal and external radiation, including from cosmic rays.




Figure 1: measured ⁸⁵Kr activities in weekly samples of air, collected at Jungfrauoch (3500 m a s l) and at Schauinsland (1000 m a s l) in the last two years.

Jungfrauoch is preferred as sampling site because there the equilibrium ^{85}Kr activity in the northern troposphere can best be determined; at this altitude admixtures of contaminated air are less probable. This equilibrium tropospheric level corresponds in Figure 1 to the lowest measured values of about 1.4 Bq/m^3 . To compensate for the yearly loss of activity in the atmosphere by radioactive decay a yearly emission rate of $4 \cdot 10^{17} \text{ Bq}$ from reprocessing plants can be estimated.

Superimposed to the basic level are irregular spikes of higher activity. This happens when air masses from reprocessing plants reach the sampling site without enough dilution with uncontaminated air. Increased activity values up to 2.7 Bq/m^3 are measured in 2005 in samples collected at the low altitude station Schauinsland, whereas at Jungfrauoch the highest value reaches “only” 1.8 Bq/m^3 (end of January 2005). Several increased values at Jungfrauoch correlate with high values in Freiburg; probably the origin of the excess ^{85}Kr is the same for both sampling sites. Forward and backward wind trajectories help to define the origin of the increased ^{85}Kr activities; usually La Hague (France) and Sellafield UK) can be distinguished. From Figure 1 it can be seen that in both years no spikes occurred in August and September. It can be concluded that emissions were lower during the summer stop of operations.

Key words:

Krypton, ^{85}Kr , radioactivity in air, reprocessing plants

Internet data bases:

HSartorius@bfs.de

Collaborating partners/networks:

purtschert@climate.unibe.ch

Scientific publications and public outreach 2005:

Umweltradioaktivität und Strahlendosen in der Schweiz, Bundesamt für Gesundheit, Abteilung Strahlenschutz, 2005 (in preparation).

Address:

Bundesamt für Strahlenschutz
Rosastrasse 9
D-79098 Freiburg

Contacts:

H. Sartorius
e-mail: HSartorius@bfs.de